

The second situation has to deal with a very different problem (problem II), which is involved for diffusion limited reactions in confined media. We consider a free diffusing reactant A that enters in a cavity, and which can react with a given fixed center i . We assume that each time the walker reaches the reactive site i , it has a probability p to react, which schematically mimics an imperfect reaction in confined geometry. Actually, numerous chemical reactions, ranging from trapping in supermolecules [24] to activation processes of synaptic receptors [25, 26] can be roughly rephrased by this generic

scheme. The question we address here, which does not seem to have been considered before, is the following: what is the probability for A to react with the center i before exiting the cavity? More generally, for a random walker starting from a site S , what is the probability Q to react with i before reaching a target site T , possibly different from S . Partitioning over the number of times the reactive site i has been visited, we have:

$$Q = 1 - \sum_{k=0}^{\infty} P(\mathbf{N}_i = k)(1-p)^k. \quad (1)$$

Once again, the random variable $\mathbf{N}_i$ is involved, but that time the determination of the *entire distribution* $P(\mathbf{N}_i = k)$ is needed.

In this Letter, we propose for the first time a method of computation of the statistics of $\mathbf{N}_i$ in confining geometry. In particular, we obtain explicitly the *exact distribution* in the case of parallelepipedic confining domains. Applications to the above mentioned examples are discussed.

We start with the computation of the mean $\langle \mathbf{N}_i \rangle$, assuming for the time being that the starting and target sites are different ($S \neq T$). We note w_{ij} the transition probabilities from site j to site i . We have $\sum_i w_{ij} = 1$, and we take $w_{ij} = w_{ji}$. These general transition probabilities can take into account reflecting boundary conditions. We consider an outgoing flux J of particles in S . Since the domain is finite, all the particles are eventually absorbed in T , and, in the stationary regime, there is an incoming flux J of particles in T . The mean particle density ρ_i thus satisfies the following equation:

$$\rho_i = \sum_j w_{ij} \rho_j + J \delta_{iS} - J \delta_{iT}, \quad (2)$$

with the boundary condition $\rho_T = 0$ (it is the absorbing site). To find the mean occupation time, we can simply notice that the mean particle density ρ_i is equal to $\langle \mathbf{N}_i \rangle J$. To solve this problem, we use the pseudo-Green function H [27, 28], which satisfies:

$$H(\mathbf{r}_i | \mathbf{r}_j) = \sum_k w_{ik} H(\mathbf{r}_k | \mathbf{r}_j) + \delta_{ij} - \frac{1}{V}, \quad (3)$$

where V is the total number of sites of the lattice. It is also symmetrical in its arguments, and the sum $\sum_i H(\mathbf{r}_i | \mathbf{r}_j)$ is a constant independent of j . Using the concise notation $H_{ij} = H(\mathbf{r}_i | \mathbf{r}_j)$, it can be seen by direct substitution that ρ_i given by

$$\langle \mathbf{N}_i \rangle = \frac{\rho_i}{J} = H_{iS} - H_{iT} + H_{TT} - H_{ST}, \quad (4)$$

satisfies Eq.(2) as well as the boundary condition $\rho_T = 0$. Note that these results also give the mean occupation time of a subdomain, which is simply the sum of the mean occupation time of all the sites in the subdomain. In particular, we can check that the mean occupation

time for the whole domain, $\sum_{i=1}^V \langle \mathbf{N}_i \rangle = V(H_{TT} - H_{ST})$, gives back the MFPT from S to T [28, 29].

Before we go further, it is necessary to give a few elements on the evaluation of H for isotropic random walks. The following exact expression [28, 30] is known in two dimensions for rectangles:

$$H(\mathbf{r} | \mathbf{r}') = \frac{4}{N} \sum_{m=1}^{X-1} \sum_{n=1}^{Y-1} \frac{\cos \frac{m\pi x'}{X} \cos \frac{n\pi y'}{Y} \cos \frac{m\pi x}{X} \cos \frac{n\pi y}{Y}}{1 - \frac{1}{2} (\cos \frac{m\pi}{X} + \cos \frac{n\pi}{Y})} + \frac{4}{N} \sum_{m=1}^{X-1} \frac{\cos \frac{m\pi x'}{X} \cos \frac{m\pi x}{X}}{1 - \cos \frac{m\pi}{X}} + \frac{4}{N} \sum_{n=1}^{Y-1} \frac{\cos \frac{n\pi y'}{Y} \cos \frac{n\pi y}{Y}}{1 - \cos \frac{n\pi}{Y}}, \quad (5)$$

where X and Y are the dimensions of the rectangle, and the coordinates x and y are half-integers going from $1/2$ to $X - 1/2$ or $Y - 1/2$. There is also a similar expression for parallelepipedic domains in three dimensions. In more general domains, the most basic approximation (which usually gives a good order of magnitude) is to approximate H by the infinite-space lattice Green function G_0 [27], G_0 being evaluated as $G_0(\mathbf{r} | \mathbf{r}') = 3/(2\pi |\mathbf{r} - \mathbf{r}'|)$ for $\mathbf{r} \neq \mathbf{r}'$, and $G_0(\mathbf{r} | \mathbf{r}) = 1.516...$ in three dimensions, and $G_0(\mathbf{r} | \mathbf{r}') = -(2/\pi) \ln |\mathbf{r} - \mathbf{r}'|$ for $\mathbf{r} \neq \mathbf{r}'$, and $G_0(\mathbf{r} | \mathbf{r}) = 1.029...$ in two dimensions. More accurate approximations can be found [28], but the above approximations are good enough to capture the qualitative behavior of the pseudo-Green function, and of the distribution of the occupation time.

It is indeed possible to obtain not only the mean, but also the entire distribution of the occupation time. The idea to tackle this *a priori* difficult problem is to use recent results concerning the so-called splitting probabilities [14, 28, 29]. In presence of two targets T_1 and T_2 , the splitting probability P_1 to reach T_1 before T_2 is [28, 29]:

$$P_1 = \frac{H_{1S} + H_{22} - H_{2S} - H_{12}}{H_{11} + H_{22} - 2H_{12}} \quad (6)$$

Denoting here $P_{ij}(i|S)$ the splitting probability to reach i before j , starting from S , we have $P(\mathbf{N}_i = 0) = P_{iT}(T|S)$, and for $k \geq 1$:

$$P(\mathbf{N}_i = k) = P_{iT}(i|S) \left[\sum_j w_{ji} P_{iT}(i|j) \right]^{k-1} \left[\sum_j w_{ji} P_{iT}(T|j) \right] \quad (7)$$

The three terms of this last equation correspond respectively to the probability to reach i before T , starting from S , the probability to return to i before reaching T , starting from i , to the power $k - 1$, and the probability to reach T before returning to i . It can thus be written

$$P(\mathbf{N}_i = k) = AB(1 - B)^{k-1} \quad \text{for } k \geq 1, \quad (8)$$

with

$$A \equiv P_{iT}(i|S) = \frac{H_{iS} + H_{TT} - H_{ST} - H_{iT}}{H_{ii} + H_{TT} - 2H_{iT}}, \quad (9)$$

and

$$B \equiv \sum_j w_{ji} P_{iT}(T|j) = 1 - \sum_j w_{ji} P_{iT}(i|j) \quad (10)$$

$$= \frac{\sum_j w_{ji} H_{Tj} - H_{iT} - \sum_j w_{ji} H_{ji} + H_{ii}}{H_{ii} + H_{TT} - 2H_{iT}} \quad (11)$$

$$= \frac{1}{H_{ii} + H_{TT} - 2H_{iT}}, \quad (12)$$

using Eq.(3), and $\sum_i w_{ij} = 1$. It can also be noted that $P(\mathbf{N}_i = 0) = 1 - A$. The distribution of the occupation numbers given by Eqs.(8)-(12) is the main result of this Letter, and several comments are in order. (i) Expressions of H given in Eq.(5) makes this result exact and completely explicit for parallelepipedic domains. (ii) Computing $\langle \mathbf{N}_i \rangle$ with this distribution gives back the expected result (4). (iii) It can be noted here that B , which characterizes the decay of the probability distribution of $\mathbf{N}_i$, is independent of the source. In addition, qualitatively, the basic evaluations of H following Eq.(5) (namely $H = G_0$) give for B the following order of magnitude, if i and T are at a distance R :

$$B \simeq \begin{cases} [2G_0(0) - 3/(\pi R)]^{-1} & \text{in 3D,} \\ [2G_0(0) + (4/\pi) \ln R]^{-1} & \text{in 2D,} \end{cases} \quad (13)$$

where $G_0(0) = G_0(\mathbf{r}|\mathbf{r})$ is a dimension-dependant constant, given in the discussion on the evaluation of H . This shows that B decreases with the distance between i and T : a larger distance corresponds to a slower decay. But, while it tends towards 0 in two dimensions (which corresponds to a wide distribution of $\mathbf{N}_i$, and a large variance), it tends to a finite value in three dimensions. It can thus be said that the sites much further from the target than the source have, in three dimensions, a significant probability to be visited, but a low probability to be visited many times, whereas, in two dimensions, they have a low probability to be visited at all, but a comparatively high probability to be visited many times. This is connected with the transient or recurrent character of the free random walk in two or three dimensions. (iv) The results obtained here for different starting and target sites may easily be adapted to identical starting and target sites ($S = T$):

$$P(\mathbf{N}_i = 0) = 1 - B; \quad P(\mathbf{N}_i = k) = B^2(1 - B)^{k-1} \quad \text{for } k \geq 1. \quad (14)$$

Note that this gives in particular a mean occupation time of 1 for all sites, a result which could be derived from an extension of Kac's formula [1, 28]. However, here, we obtain not only the mean occupation number but the entire distribution of this occupation number, which appears to vary from site to site: the further the site is from the target, the slower the probability distribution decays.

We now discuss the applications of these general results to the examples mentioned in the introduction. As for the random trap model (problem I), we focus here on the especially interesting case of a one-sided Levy

stable distribution [2] $\psi(t) = f_\alpha(t, \tau_0 \alpha \cos(\pi\alpha/2), 1, 0)$ ($0 < \alpha < 1$), which corresponds to an algebraic decay:

$$\psi(t) \sim \frac{\alpha \tau_0^\alpha}{\Gamma(1 - \alpha) t^{1+\alpha}} \quad (15)$$

and whose Laplace transform is $\hat{\psi}(u) = \exp(-\tau_0^\alpha u^\alpha)$ (τ_0 can be seen as the typical waiting time). The Laplace transform $\hat{\pi}(u)$ of the distribution of the MFPT with respect to the disorder reads

$$\hat{\pi}(u) = \prod_{i=1}^V \hat{\psi}(\langle \mathbf{N}_i \rangle u) = \exp(-(T_{typ} u)^\alpha) \quad (16)$$

The probability density of the MFPT is then as could have been expected a one-sided Levy stable law, but with a non trivial typical time:

$$T_{typ} = \tau_0 \left(\sum_{i=1}^V (H_{iS} - H_{iT} + H_{TT} - H_{ST})^\alpha \right)^{1/\alpha} \quad (17)$$

For large size domain V , this result can be applied to any wide-tailed distribution of the waiting times satisfying Eq.(15) [12]. It can be shown that T_{typ} is bounded by $\tau_0 V^{1/\alpha} (H_{TT} - H_{ST})$, and tends towards this upper bound as V grows, which provides a simple estimation of T_{typ} and indicates that for large enough domains, the scaling of T_{typ} with the source and target positions is the same as for the discrete-time random walk (pure systems) [28, 29]. We thus showed that the random trap problem in confined geometries, with a wide-tailed waiting time distribution, has a Levy distribution of mean first-passage times, with a non-trivial typical time. The scaling with the size V is $V^{1/\alpha}$. The scaling with the source and target positions is modified by the disorder in small confining domains, while it is the same as for pure systems in large enough domains.

Concerning the application to diffusion-limited reactions (problem II), the probability Q to have reacted with i before reaching T writes, using Eqs.(1),(8):

$$Q = \frac{Ap}{1 - (1 - p)(1 - B)} \quad (18)$$

The expression (18) displays a subtle interplay between the geometrical factors, involved through the terms A and B , and the reactivity p . Focusing now on the specific case of identical starting and target points (meaning $A = B$, cf. Eq.(14)), we exhibit two interesting limiting regimes. In the ‘‘reactivity limited regime’’, defined by $p \ll B$, we have $Q \sim p$. In particular, in that regime Q does not depend on the reactive site i . In other words, for a fixed reactivity p , all sites i such that $p \ll B$ have the same probability of reaction Q , and the detailed position of i does not come into play. On the contrary, the ‘‘geometrically limited regime’’ $p \gg B$ leads to $Q \sim B$, which does no longer depend on p , but only on the geometry. Given the order of magnitude of B (cf.




FIG. 2: Color online. Simulations (symbols) versus analytical prediction (lines) Eq.(18) of the probability Q to react before returning to the target site as a function of the probability to react at site i . The confining domain is a square of side 51, and the target is at the middle of an edge, of coordinates (0,25), the site (0,0) being a corner site. The three curves corresponds to different positions of site i : (1,25) (red upper curve), (25,25) (blue mid curve), and (50,25) (green lower curve).

Eq. (13)), this can essentially happen in two dimensions, when $R \gg \exp(-\pi/(2p))$. This can be explained by the recurrent character of the two-dimensional random walk: when the reacting site i is far enough from the target, if a random walker reaches it, it is likely to visit it many times before returning to T , and is thus almost sure to react whenever i is reached. The reaction probability Q then becomes the probability to reach the site i . Consequently, the position of the reacting site has a low influence on reactivity in three dimensions, or when the reacting site is within a disk of radius $R = \exp(-\pi/(2p))$ around the target in two dimensions. If the reacting site is further, the

geometrical effects become preeminent. We show in Fig. 2 a graph of Q , as a function of p , for different positions of i (near the target, in the middle of the domain, and at the opposite), the source and target point being identical. The limiting regimes can be well identified.

To conclude, we have computed the distribution of the occupation time of a given site i , for a random walk in confined geometry, eventually trapped at a target. This distribution is exact, and completely explicit in the case of parallelepipedic confining domains. While the mean occupation time, unsurprisingly, is higher when i is near the source and lower near the target (and uniform if the source and target are identical), the distribution of the occupation time is essentially exponential, with a slower decay when the point is far away from the target. We have also presented important applications of these results in two different fields. The first one is transport in quenched disorder media: The mean first passage time for the random trap model has been computed for the first time in dimensions greater than 1, and has been shown to display a non-trivial dependence with the source and target positions. The second application is to diffusion limited reactions in confined geometry: The probability of reaction with a given imperfect center before being trapped by another one has been explicitly calculated, and has proven to present a complex dependence both in the geometrical and chemical parameters. We believe that the results obtained in this Letter could be relevant to systems involving diffusion in confining domains, displaying inhomogeneous physical or chemical properties. .

-
- [1] D. Aldous and J. Fill, *Reversible markov chains and random walks on graphs* (<http://www.stat.berkeley.edu/~aldous/RWG/book.html>, 1999).
 - [2] B. Hughes, *Random walks and random environments* (New York: Oxford University Press, 1995).
 - [3] C. Godrèche and J. Luck, J. Stat. Phys. **104**, 489 (2001).
 - [4] S. Majumdar and A. Comtet, Phys. Rev. Lett. **89**, 060601 (2002).
 - [5] O. Bénichou, M. Coppey, J. Klafter, M. Moreau, and G. Oshanin, J. Phys. A **36**, 7225 (2003).
 - [6] S. Blanco and R. Fournier, Europhys. Lett. **61**, 168 (2003).
 - [7] O. Bénichou, M. Coppey, M. Moreau, P. Suet, and R. Voituriez, Europhys. Lett. **70**, 42 (2005).
 - [8] S. Condamin, O. Bénichou, and M. Moreau, Phys. Rev. E **72**, 016127 (2005).
 - [9] S. Blanco and R. Fournier, Phys. Rev. Lett. **97**, 230604 (2006).
 - [10] S. Burov and E. Barkai, Phys. Rev. Lett. **98**, 250601 (2007).
 - [11] M. Ferraro and L. Zaninetti, Physica A **338**, 307 (2004).
 - [12] J-P. Bouchaud and A. Georges, Phys. Rep. **195**, 127 (1990).
 - [13] O. Bénichou, M. Coppey, M. Moreau, and G. Oshanin, J. Chem. Phys. **123**, 194506 (2005).
 - [14] S. Redner, *A guide to First-Passage Processes* (Cambridge University Press, 2001).
 - [15] P. Levitz, D. Grebenkov, M. Zinsmeister, K.M. Kolwankar, and B. Sapoval, Phys. Rev. Lett. **96**, 180601 (2006).
 - [16] A. Blumen, J. Klafter, and G. Zumofen, Phys. Rev. B **28**, 6112 (1983).
 - [17] S.B. Yuste and K. Lindenberg, Chem. Phys. **284**, 169 (2002).
 - [18] S. Yuste and L. Acedo, Phys. Rev. E **64**, 061107 (2001).
 - [19] S. Yuste and L. Acedo, Phys. Rev. E **68**, 036134 (2003).
 - [20] O. Bénichou, M. Coppey, M. Moreau, P. Suet, and R. Voituriez, Phys. Rev. Lett. **94**, 198101 (2005).
 - [21] K. Murthy and K. Kehr, Phys. Rev. A **40**, 2082 (1989).
 - [22] E. Hernandez-Garcia and M. Caceres, Phys. Rev. A **42**, 4503 (1990).
 - [23] M. Kawasaki, T. Odagaki and K.W. Kehr, Phys. Rev. B **67**, 134203 (2003).
 - [24] A. Bar-Haim and J. Klafter, J. Chem. Phys. **109**, 5187 (1998).
 - [25] L. Savtchenko and D. Rusakov, PNAS **104**, 1823 (2007).
 - [26] D. Holcman, A. Marchewka, and Z. Schuss, Phys. Rev. E **72**, 031910 (2005).
 - [27] G. Barton, *Elements of Green's Functions and Propaga-*

- tion* (Oxford Science Publications, 1989).
- [28] S. Condamin, O. Bénichou, and M. Moreau, Phys.Rev.E **75** (2007).
- [29] S. Condamin, O. Bénichou, and M. Moreau, Phys. Rev. Lett. **95**, 260601 (2005).
- [30] S. Condamin and O. Bénichou, J.Chem.Phys. **124**, 206103 (2006).